

Determination of the Critical Point and Exponents from Short-time Dynamics

L. Schülke and B. Zheng

Universität – GH Siegen, D – 57068 Siegen, Germany

Abstract

The dynamic process for the two dimensional three state Potts model in the critical domain is simulated by the Monte Carlo method. It is shown that the critical point can rigorously be located from the universal short-time behaviour. This makes it possible to investigate critical dynamics independently of the equilibrium state. From the power law behaviour of the magnetization the exponents $\beta/(\nu z)$ and $1/(\nu z)$ are determined.

1 Introduction

Recently more and more attention is drawn to the universal behaviour of the critical dynamic process in the macroscopic short-time regime. for the critical dynamic process of model A for the $O(N)$ vector model starting from a random state with a small but non-zero initial magnetization it has been argued with ϵ -expansion up to two loop that the magnetization undergoes a *critical initial increase* [1]. Such a scenario has been simulated for the Ising model and the Potts model by the Monte Carlo method [2, 3], and the new dynamic exponent θ which governs the initial universal behaviour has directly been determined. The exponent θ may also be estimated indirectly from the power law decay of the auto-correlation [4, 5], and with the damage spreading technique [6]. Universality [7, 8] as well as a generalized finite size scaling [2, 9, 10] has also been confirmed. Such an investigation of the short-time critical dynamics not only enlarges our fundamental knowledge on critical phenomena but also provides possible improvement on the numerical determination of the dynamic and the static exponents. The generalization and application to complex systems are also undergoing [4, 11, 12, 13].

Of special interest is here a previous paper by the authors where the short-time dynamics for the Potts model at the critical point was simulated with the Monte Carlo Heat-bath algorithm [3]. We have concentrated ourselves on the critical initial increase of the magnetization as well as the power law behaviour of the auto-correlation and the second moment of the magnetization. The new dynamic exponent θ has directly been measured and the traditional dynamic exponent z as well as β/ν were also obtained. The results relate the direct and indirect measurement of θ to each other and provide confident support for the dynamic scaling in the short-time regime. Especially the measured dynamic exponent z is quite rigorous. Compared with the distributed values of z between $z = 2.2$ and $z = 2.7$ from different simulations [14, 15, 16, 17], the exponent z we obtained supports the small one [14, 15].

However some important problems remain open in the numerical simulation of the short-time dynamics. First, the critical point was *always input* in the numerical investigation of the short-time dynamics. The critical point is usually defined and measured in the equilibrium state of the systems.. It is be very interesting to know whether one can define and locate the critical point from the short-time dynamics itself or not. This would allow a complete investigation of the short-time dynamics independent of the equilibrium state. On the other hand, for the critical system (with infinite volume) exactly at the critical point the dynamic process will, in principle, never reach the equilibrium state due to the critical slowing down, at least for the *local dynamics*. Therefore it is also important from the practical point of view that one can determine the critical point from the dynamics itself. Furthermore one may imagine that the critical point for the dynamic system could be different from that for the equilibrium state. This may conceptually broaden the scope of the critical phenomena.

Actually some discussion of this kind has been made with respect to the damage spreading [18]. One purpose of this letter is to clarify how to locate rigorously the critical point from the short-time dynamics.

Secondly, how to determine the static exponent β and ν from the short-time dynamics in a rigorous way still requires further investigation. It has been pointed out that the determination of the static exponents from the short-time dynamics is quite promising since one does not enter the long-time regime of the dynamic evolution where the critical slowing down is severe [9, 10]. It has been demonstrated for the Ising model that the static exponents β and ν may be estimated from finite size scaling. One can easily realize that the measurement of the static exponents from the dynamics depends essentially on the quality of the measurement of the dynamic exponent z . In some cases a small relative error for z may induce big errors for the static exponents. To determine the dynamic exponent z independently with the time-dependent Binder cumulant in principle works well. But it happens sometimes that the cumulant is somewhat fluctuating since it is constructed from higher moments of the magnetization.

Therefore another purpose of this Letter is to provide an alternative way to determine the static exponents β and ν from the power law behaviour of the observables in the beginning of the time evolution of the dynamic process. Compared with the method based on finite size scaling, the measurement of the exponents from the power law behaviour of the observables is more direct. The exponents may be extracted from a single lattice and the finite size effect may easily be under control. Actually in reference [3] the static exponent β/ν has been obtained from the power law increase of the second moment. However the result is still rough. What one practically measures from the second moment of the magnetization is $(d - 2\beta/\nu)/z$. As it was pointed out [3], the value of β/ν is much smaller than d and z , and therefore a small relative error for z or $(d - 2\beta/\nu)/z$ will induce a big error for $2\beta/\nu$. We should search for a better way.

Let us start with the determination of the critical point and the exponent $1/(\nu z)$. Following the consideration in [3], we first study the critical dynamic process of model A for the two dimensional three state Potts model starting from a random initial state with small but non-zero initial magnetization. The Hamiltonian is given by

$$\mathcal{H} = J \sum_{\langle i,j \rangle} \delta_{\sigma_i \sigma_j}; \quad \sigma_i = -1, 0, 1. \quad (1)$$

According to the argument of Janssen, Schaub and Schmittmann obtained with the renormalization group method, one may expect a generalized scaling relation

$$M^{(k)}(t, \tau, m_0) = b^{-k\beta/\nu} M^{(k)}(b^{-z}t, b^{1/\nu}\tau, b^{x_0}m_0). \quad (2)$$

to be valid also for the Potts model. for the magnetization ($k = 1$), taking $b = t^{1/z}$ and assuming $t^{x_0/z}m_0$ is small enough, we can expand with respect to

$t^{x_0/z}m_0$ and obtain

$$M(t, \tau, m_0) = m_0 t^\theta F(t^{1/(\nu z)}\tau) \quad (3)$$

where $\theta = (x_0 - \beta/\nu)/z$. It is clear that for $\tau = 0$ we observe the so-called critical initial (power law) increase of the magnetization. for $\tau \neq 0$, but still in the critical domain the time evolution of the magnetization will be modified by the scaling function $F(t^{1/(\nu z)}\tau)$. Such a fact can be used for the determination of the critical point. In Fig. 1 the time evolution of the magnetization for three different coupling $J = 1.0025$, $J = 1.0050$, and $J = 1.0075$, with initial magnetization $m_0 = 0.04$ and lattice size $L = 72$ are displayed with solid lines in double-log scale. The coupling for the line in the centre, $J = 1.0050$, corresponds to the theoretical critical point for the infinite system. The total number of samples with independent initial configurations for each coupling J is 600 000. By a linear extrapolation in J between the simulated curves one can find a particular coupling which gives the best power law behaviour for the magnetization. This is the “critical point” for the finite system. In principle there can also be some effect from the finite initial magnetization m_0 . The dots in Fig. 1 represent the time evolution of the magnetization at such a critical point $J_c = 1.0055(8)$. It is remarkable that the critical point located in this way is so close to the exact one. To estimate the critical point, we have linearly interpolated the magnetization $M(t, J)$ between the measured curves for $J = 1.0025$ and $J = 1.0075$. At the time t , the slope is measured within a time interval $[t, t + 15]$. Theoretically, at the critical point J_c , the slope $\theta(t, J_c)$ is constant with respect to t . Practically the critical point is determined by searching for the minimum square deviation of the slope $\theta(t, J)$ from a constant. In Fig. 2 this deviation is shown when the coupling is varying around the critical value. The clear minimum is a confident indication for the critical point. It has been pointed out that in the measurement of the exponent θ , the microscopic time scale t_{mic} for the Heat-Bath algorithm is ignorably small. Therefore in the above estimate of the critical point, the time interval is taken to be $[1, 100]$. Actually the time interval may be taken even smaller, e.g. $[1, 50]$. This is really the physics from the *short-time regime* of the dynamic evolution. Furthermore, examining the result for smaller lattice size $L = 32$ and somewhat bigger initial magnetization $m_0 = 0.08$, one can realize that the effects of finite size and m_0 for $L = 72$ and $m_0 = 0.04$ are already very small, at least in the determination of the critical point. This is an advantage for determining the critical point from the short-time dynamics.

Theoretically the critical point could be located from short-time behaviour of any observable in the critical domain. However practically many of them are not very sensitive to the coupling. Also to determine the critical point efficiently the dependence of the chosen observable on the critical exponents should be as small as possible. For example, the second moment of the magnetization and the auto-correlation are not good choices for the determination of the critical point. In reference [2] it was mentioned that one might be able to locate the

critical point from the time-dependent cumulant. But this also appears to have some complication since higher moments are involved.

To determine the exponent $1/(\nu z)$, we differentiate $\ln M(t, \tau, m_0)$ with respect to τ at the critical point, and for small enough $t^{x_0/z} m_0$

$$\partial_\tau \ln M(t, \tau, m_0)|_{\tau=0} = t^{1/(\nu z)} \partial_\tau \ln F(\tau')|_{\tau'=0} \quad (4)$$

Therefore $\partial_\tau \ln M(t, \tau, m_0)|_{\tau=0}$ also shows a power law increase. In the numerical simulation we have approximated the differentiation by the difference between $J = 1.0025$ and $J = 1.0075$. This derivative is plotted in Fig. 3 in double-log scale. From the slope one can estimate the critical exponent $1/(\nu z)$. Using the exponent measured from the auto-correlation as input, one obtains the static exponent $1/\nu$. The result together with the exponent z and the measured critical point is given in table 1. Due to some possible effect of t_{mic} , we have made the measurement within the time interval $[20, 100]$. Compared with the exact value $1/\nu = 1.2$ the measured value is slightly bigger. This comes probably from the approximation of the differentiation by the difference. To improve the quality of $1/\nu$ we might represent alternatively the derivative by higher moments [19]. Here we should point out that the slight difference between the values for z in table 1 and that in reference [3] comes from some more serious consideration of the effect of t_{mic} and the finite lattice size [8].

J_c	z	β/ν	$1/\nu$
1.0055(8)	2.196(8)	0.1330(3)	1.24(3)

Table 1: Results for J_c , β/ν and $1/\nu$ obtained in this paper. The value for z used as an input has been taken from the previous investigation.

Now we turn to the determination of the static exponent $\beta/(\nu z)$. It has been discussed before that it is not the best way to measure it from the second moment. Due to the very short spatial correlation length in the very beginning of the time evolution of the dynamic process the second moment behaves as $M^{(2)}(t, L) \sim L^{-d}$. Therefore obtaining a good power law behaviour for the second moment is not so easy, especially for relative big lattices in the short-time regime. Fortunately recent investigations of the dynamic Ising model [20, 10] shows that from the power law *decay* of the magnetization

$$M(t) \sim t^{-\frac{\beta}{\nu z}} \quad (5)$$

in the dynamic process starting from a *completely ordered state*, i.e. $m_0 = 1$, it is possible to estimate the exponent $\beta/(\nu z)$. With z in hand, we can obtain the static exponent β/ν . Here we have simulated such a dynamic process for the Potts model. We have pointed out before that the critical point we estimated is

very close to the exact one. Therefore in the simulation here we take the exact value for the critical point for simplicity. The results support that universality and scaling appear to be valid also in a quite early stage of the time evolution and one can really obtain the exponent $\beta/(\nu z)$ quite accurately in the short-time dynamics. In Fig. 4 the time evolution of the magnetization is displayed for lattice sizes $L = 72$ and $L = 144$ in double-log scale. The total number of samples for independent initial configurations is 200000 for $L = 72$ and 80000 for $L = 144$. One can hardly see any difference between both curves, i.e., there is already an invisible finite size effect for lattice size $L = 72$. From the slope of the curves one can estimate the exponent $\beta/(\nu z)$. Detailed analysis reveals that in this case $t_{mic} \sim 50$, which is bigger than that in the measurement of θ . However the behaviour becomes very stable after t_{mic} . Measuring the slope in a time interval $[100, 250]$ and using the value of z as input, one obtains the static exponent $\beta/\nu = 0.1326(6)$ for $L = 72$ and $\beta/\nu = 0.1333(2)$ for $L = 144$. The errors are estimated by dividing the samples into two or six groups. The average value is given in table 1. The static exponent β/ν obtained in this way is very rigorous. In general the quality of the measurement for the first moment is better than that of the second moment. Besides the magnetization in this process is strongly self-averaging and therefore one can easily simulate bigger lattices. Furthermore the error of the input z will now have almost no effect on the quality of the exponent β/ν since the value of $\beta/(\nu z)$ is much smaller than that of z . This situation is very different from that for the measurement of the second moment.

In conclusion, we have numerically simulated the critical dynamic process of model A for the two dimensional three state Potts model starting from both a random state with small initial magnetization and a completely ordered state. From the universal power law behaviour of the magnetization in the short-time regime of the dynamic evolution, the critical point and the static exponents are obtained. The results provide further confirmation for the dynamic scaling in the short-time dynamics, including the extension to the dynamic process starting from a completely ordered state. Especially the measurement of the critical point and the exponent β/ν is very rigorous. This makes it possible to study the short-time dynamics completely independent of the equilibrium state. The determination of the static properties from the short-time dynamics is not only conceptually interesting but also practically applicable. The application of this technique to the unsolved critical systems is being carried out.

References

- [1] H. K. Janssen, B. Schaub and B. Schmittmann. *Z. Phys.* **B 73** 539 (1989).
- [2] Z. B. Li, U. Ritschel and B. Zheng. *J. Phys. A: Math. Gen.* **27** L837 (1994).
- [3] L. Schülke and B. Zheng. *Phys. Lett.* **A 204** 295 (1995).

- [4] D. A. Huse. *Phys. Rev.* **B 40** 304 (1989).
- [5] K. Humayun and A. J. Bray. *J. Phys. A* **24** 1915 (1991).
- [6] P. Grassberger. *Physica A* **214** 547 (1995).
- [7] N. Menyhárd. *J. Phys. A: Math. Gen.* **27** 663 (1994).
- [8] K. Okano, L. Schülke, K. Yamagishi and B. Zheng. Universality and Scaling in Short-Time Critical Dynamics. *preprint* SI-95-7. Univ. of Siegen.
- [9] Z. B. Li, L. Schülke and B. Zheng. *Phys. Rev. Lett.* **74** 3396 (1995).
- [10] Z. B. Li, L. Schülke and B. Zheng. Finite Size Scaling and Critical Exponents in Critical Relaxation. *Phys. Rev. E*. to be published.
- [11] S. J. Lee, J. R. Lee and B. Kim. *Phys. Rev.* **E 51** R4 (1995).
- [12] A. D. Rutenberg and A. J. Bray. *Phys. Rev.* **E 51** R1641 (1995).
- [13] R. E. Blundell, K. Humayun and A. J. Bray. *J. Phys. A, Math. Gen.* **25** L733 (1992).
- [14] M. Aydin and M. C. Yalabik. *J. Phys. A, Math. Gen.* **18** 1741 (1985).
- [15] S. Tang and D. P. Landau. *Phys. Rev.* **B 36** 567 (1987).
- [16] K. Binder. *Z. Phys.* **B 43** 119 (1981).
- [17] M. Aydin and M. C. Yalabik. *J. Phys. A, Math. Gen.* **21** 769 (1988).
- [18] P. Grassberger. *J. Phys. A, Math. Gen.* **28** L67 (1995).
- [19] A. M. Ferrenberg and D. P. Landau. *Phys. Rev.* **B 44** 5081 (1991).
- [20] D. Stauffer. *Physica A* **186** 197 (1992).




Figure 1: The magnetization $M(t)$ for initial magnetization $m_0 = 0.04$ plotted versus time in a double-log scale. The solid lines correspond to three different values for J , $J = 1.0025$, $J = 1.0050$ and $J = 1.0075$. The dotted curve represents the magnetization for the critical point $J_c = 1.0055$ which shows the best power law behaviour.




Figure 2: Deviation from the power law behaviour when J is varied around the critical point.




Figure 3: The derivative $\partial_\tau \ln M(t, \tau, m_0)|_{\tau=0}$ plotted versus time in a double-log scale. In order to avoid possible effects of t_{mic} , the slope has been calculated from the time interval $t = [20, 100]$. The dotted curve represents a straight line with the measured slope.




Figure 4: The magnetization $M(t)$ versus time in a double-log scale. The solid line and the crosses on that line represent the results for $L = 72$ and $L = 144$. The slope has been obtained from the time interval $t = [100, 250]$ in order to avoid effects from t_{mic} . The dotted curve is the straight line with the measured slope.